

Ibuprofen (Motrin, Advil) Dosage Chart

Give the weight-appropriate dosage listed below every 6 - 8 hours as needed for temperature over 100.4°F or for pain. Always give with food. **Do not exceed 4 doses in a 24-hour period.**

Do not give Ibuprofen if child is less than 6 months old or if your child has a kidney problem

Weight (pounds)	Infant Drops 50mg / 1.25ml	Childrens Liquid 100mg / 5ml
8-9	40 mg (1.0 ml)	
10-11	50 mg (1.25 ml)	
12-13	60 mg (1.5 ml)	
14-16	70 mg (1.75 ml)	
17-18	80 mg (2.0 ml)	
19-22	90 mg (2.25 ml)	
22-26	100 mg (2.5 ml)	5 ml (1 tsp)
28-31	120 mg (3.0 ml)	6 ml
32-35	140 mg (3.5 ml)	7 ml (1.5 tsp)
36-39	160 mg (4.0 ml)	8 ml
40-44		9 ml
4 5- 4 8		10 ml (2 tsp)
49-53		11 ml
54-57		12 ml
58-62		13 ml
63-66		14 ml
67-71		15 ml (3 tsp)
72-75		16 ml
76-80		17 ml (3.5 tsp)
81-84		18 ml
85-88		19 ml
89 lbs and up		20 ml (4 tsp)

Tylenol (acetaminophen) Dosage Chart

Give the weight-appropriate dosage listed below every 4-6 hours as needed for a temperature over 100.4 F. Do not exceed 5 doses in a 24-hour period.

Do not give Tylenol to children under 2 months of age or if they have not had their first set of immunizations without first consulting a doctor. Do not use Tylenol if your child has any liver problems.

Infants' Acetaminophen Oral Suspension 160 mg / 5 ml		Chewable Tablets 80 mg tablets	Junior Strength 160 mg tablets	
Weight (lbs)	Age (mos)	Dose (ml)	Dose (tablets)	Dose (tabs or caps)
6 - 11	0-3	1.25 ml		
12-17	4-11	2.5 ml		
18-23	12-23	3.75 ml		
24-35	24-36	5 ml	2 tabs	1 tab/cap
36-39		8 ml	3 tabs	1.5 tabs/caps
40-44		9 ml	3 tabs	1.5 tabs/caps
45-48		10 ml (2 tsp)	4 tabs	2 tabs/caps
49-53		11 ml	4 tabs	2 tabs/caps
54-57		12 ml (2.5 tsp)	5 tabs	2.5 tabs/caps
58-62		13 ml	5 tabs	2.5 tabs/caps
63-66		14 ml	5 tabs	2.5 tabs/caps
67-71		15 ml (3†sp)	6 tabs	3 tabs/caps
72-75		16 ml	6 tabs	3 tabs/caps
76-80		17 ml (3.5 tsp)	7 tabs	3.5 tabs/caps
81-84		18 ml	7 tabs	3.5 tabs/caps
85-88		19 ml	7 tabs	3.5 tabs/caps
89 lbs and up		20 ml (4tsp)	8 tabs	4tabs/caps